

Representing the Mean Residual Life in Terms of the Failure Rate

R. C. GUPTA AND D. M. BRADLEY

Department of Mathematics and Statistics, University of Maine

5752 Neville Hall, Orono, ME 04469-5752, U.S.A.

rcgupta@maine.maine.edu

bradley@math.umaine.edu

dbradley@member.ams.org

(Received August 2002; accepted February 2003)

Abstract—In survival or reliability studies, the mean residual life or life expectancy is an important characteristic of the model. Whereas the failure rate can be expressed quite simply in terms of the mean residual life and its derivative, the inverse problem—namely that of expressing the mean residual life in terms of the failure rate—typically involves an integral of a complicated expression. In this paper, we obtain simple expressions for the mean residual life in terms of the failure rate for certain classes of distributions which subsume many of the standard cases. Several results in the literature can be obtained using our approach. Additionally, we develop an expansion for the mean residual life in terms of Gaussian probability functions for a broad class of ultimately increasing failure rate distributions. Some examples are provided to illustrate the procedure. © 2003 Elsevier Science Ltd. All rights reserved.

Keywords—Pearson family of distributions, Gaussian probability function, Ultimately increasing failure rate.

1. INTRODUCTION

In life testing situations, the expected additional lifetime given that a component has survived until time t is a function of t , called the mean residual life. More specifically, if the random variable X represents the life of a component, then the mean residual life is given by $m(t) = E(X - t \mid X > t)$. The mean residual life has been employed in life length studies by various authors, e.g., Bryson and Siddiqui [1], Hollander and Proschan [2], and Muth [3]. Limiting properties of the mean residual life have been studied by Meilijson [4], Balkema and de Hann [5], and more recently by Bradley and Gupta [6]. A smooth estimator of the mean residual life is given by Chaubey and Sen [7].

It is well known that the failure rate can be expressed quite simply in terms of the mean residual life and its derivative, see (2.4). However, the inverse problem—namely that of expressing the mean residual life in terms of the failure rate—typically involves an integral of a complicated expression. In this paper, we obtain a simple expression for the mean residual life in terms of the failure rate for certain classes of distributions. Many of the standard cases are subsumed, and several results in the literature can be obtained using our approach. However, the emphasis

here is to express the mean residual life in terms of the failure rate. For the class of ultimately increasing failure rate distributions, we also provide sufficient conditions under which the mean residual life can be expanded in terms of Gaussian probability functions. Finally, some examples are presented to illustrate the procedure.

2. BACKGROUND AND DEFINITIONS

Let $F : [0, \infty) \rightarrow [0, \infty)$ be a nondecreasing, right continuous function with $F(0) = 0$, $\lim_{x \rightarrow \infty} F(x) = 1$, and let ν denote the induced Lebesgue-Stieljes measure. (Equivalently, let ν be a probability measure on $[0, \infty)$ and let F be the cumulative distribution function of ν .) If X is a nonnegative random variable representing the life of a component having distribution function F , the mean residual life is defined by

$$m(t) = E(X - t | X > t) = \frac{1}{\bar{F}(t)} \int_t^\infty (x - t) d\nu(x), \quad t \geq 0,$$

where $\bar{F} = 1 - F$ is the so-called survival function. Writing $x - t = \int_t^x du$ and employing Tonelli's theorem yields the equivalent formula

$$m(t) = \frac{1}{\bar{F}(t)} \int_t^\infty \int_t^x du d\nu(x) = \frac{1}{\bar{F}(t)} \int_t^\infty \int_u^\infty d\nu(x) du = \frac{1}{\bar{F}(t)} \int_t^\infty \bar{F}(u) du, \quad (2.1)$$

which is sometimes also used as a definition. The cumulative hazard function may be defined by $R = -\log \bar{F}$. Then (2.1) implies that

$$m(t) = \int_0^\infty \exp \{R(t) - R(t + x)\} dx. \quad (2.2)$$

If F (equivalently, ν) is also absolutely continuous, then the probability density function f and the failure rate (hazard function) r are defined almost everywhere by $f = F'$ and $r = f/\bar{F} = R'$, respectively, and then

$$R(t) = \int_0^t \frac{1}{\bar{F}(x)} d\nu(x) = -\log \bar{F}(t) = -\int_0^t \frac{d\bar{F}(x)}{\bar{F}(x)} = \int_0^t r(x) dx. \quad (2.3)$$

In view of (2.2) and (2.3), we have expressed m in terms of r , albeit somewhat indirectly.

Ideally, we would like to express the mean residual life in terms of known functions of the failure rate and its derivatives without the use of integrals. In any case, it is useful to have alternative representations of the mean residual life. We note that the converse problem, that of expressing the failure rate in terms of the mean residual life and its derivatives is trivial, for (2.1) and (2.3) imply that

$$m'(t) = r(t)m(t) - 1. \quad (2.4)$$

3. A GENERAL FAMILY OF DISTRIBUTIONS, INCLUDING THE PEARSON FAMILY

Consider the family of distributions whose probability density function f is differentiable. Write

$$\frac{f'(x)}{f(x)} = \frac{\mu - x}{g(x)} - \frac{g'(x)}{g(x)}, \quad (3.1)$$

where μ is a constant and g satisfies the first-order linear differential equation,

$$g'(x) + \frac{f'(x)}{f(x)}g(x) = \mu - x.$$

Alternatively, we may view g as given, then f is uniquely determined by (3.1). Clearing the fractions in (3.1) and integrating yields

$$\int_t^\infty xf(x) dx = \mu\bar{F}(t) + g(t)f(t), \quad t \geq 0.$$

In other words,

$$E(X | X > t) = \mu + g(t)r(t), \quad t \geq 0, \quad (3.2)$$

or equivalently,

$$m(t) = \mu - t + g(t)r(t), \quad t \geq 0. \quad (3.3)$$

Thus, the mean residual life has been expressed in terms of the failure rate r , the given function g , and the constant μ . By appropriately specializing g in (3.3), one can obtain many of the important cases that have appeared in the literature. We note that a result similar to (3.3) was obtained by Ruiz and Navarro [8], but their emphasis was different.

3.1. The Pearson Family

The quadratic function $g(x) = a_0 + a_1x + a_2x^2$ with $a_2 \neq -1/2$ yields the Pearson family, special cases of which include the beta distributions, the gamma distribution, and the normal distribution. From (3.1), we have

$$\frac{f'(x)}{f(x)} = \frac{\mu - x}{a_0 + a_1x + a_2x^2} - \frac{a_1 + 2a_2x}{a_0 + a_1x + a_2x^2} = -\frac{x + d}{A_0 + A_1x + A_2x^2}, \quad (3.4)$$

say, where $A_j = a_j/(1 + 2a_2)$ for $j = 0, 1, 2$ and $d = (a_1 - \mu)/(1 + 2a_2)$. For the Pearson family, (3.2) gives

$$E(X | X > t) = \frac{A_1 - d}{1 - 2A_2} + \frac{A_0 + A_1t + A_2t^2}{1 - 2A_2}r(t), \quad (3.5)$$

see [8,9]. Note that since $A_2 = a_2/(1 + 2a_2)$, it is impossible for $1 - 2A_2$ to vanish in (3.5).

3.1.1. The beta distributions

With $g(x) = x(1 - x)/(a + b)$, we have the beta distribution of the first kind:

$$f(x) = \frac{\Gamma(a+b)}{\Gamma(a)\Gamma(b)}x^{a-1}(1-x)^{b-1}, \quad 0 < x < 1, \quad a > 0, \quad b > 0.$$

It satisfies

$$\frac{f'(x)}{f(x)} = -\frac{x + (1-a)/(a+b-2)}{x(1-x)/(a+b-2)}, \quad a+b \neq 2,$$

and hence, belongs to the Pearson family given by (3.4) with $d = (1-a)/(a+b-2)$, $A_0 = 0$, $A_1 = -A_2 = 1/(a+b-2)$, and $a+b \neq 2$. Thus,

$$E(X | X > t) = \mu + \frac{t(1-t)}{a+b}r(t), \quad 0 < t < 1,$$

where $\mu = a/(a+b)$. If $a+b = 2$, we have

$$f(x) = \frac{1}{\Gamma(a)\Gamma(2-a)} \left(\frac{x}{1-x} \right)^{a-1}, \quad 0 < a < 2,$$

and $g(x) = x(1-x)/2$, $\mu = a/2$, from which (3.2) gives

$$E(X | X > t) = \frac{1}{2}a + \frac{1}{2}t(1-t)r(t), \quad 0 < t < 1.$$

The beta distribution of the second kind has the form

$$f(x) = \frac{cx^{\beta-1}}{(\gamma+x)^{\alpha+\beta}}, \quad c, \alpha, \beta, \gamma, x > 0.$$

Since then

$$\frac{f'(x)}{f(x)} = -\frac{x + (1-\beta)\gamma/(\alpha+1)}{x^2/(\alpha+1) + \gamma x/(\alpha+1)},$$

f belongs to the Pearson family (3.4) with $d = (1-\beta)\gamma/(1+\alpha)$, $A_0 = 0$, $A_1 = \gamma/(1+\alpha)$, and $A_2 = 1/(1+\alpha)$. Hence,

$$E(X | X > t) = \mu + \left(\frac{t^2 + \gamma t}{\alpha - 1}\right) r(t), \quad \alpha \neq 1,$$

where $\mu = \beta\gamma/(\alpha-1)$ is the mean. A similar result was obtained by Ahmed [10] using a completely different approach.

3.1.2. The gamma distribution

Let $B > 0$. The linear function $g(x) = Bx$ for $x > 0$ yields the gamma distribution. In this case, (3.1) gives

$$\frac{f'(x)}{f(x)} = -\frac{x + B - \mu}{Bx},$$

so that f belongs to the Pearson family (3.4) with $d = B - \mu$, $A_0 = 0$, $A_1 = B$, and $A_2 = 0$. In fact,

$$f(x) = \frac{B^{-\mu/B}}{\Gamma(\mu/B)} x^{\mu/B-1} e^{-x/B},$$

a gamma distribution with mean μ . Hence, (3.2) takes the form

$$E(X | X > t) = \mu + Btr(t)$$

in this case. This is Theorem 1 of [11], see also [12].

3.1.3. The normal distribution

Let $\sigma > 0$. The constant function $g(x) = \sigma^2$ for $-\infty < x < \infty$ yields the normal distribution. In this case, (3.3) gives

$$\frac{f'(x)}{f(x)} = \frac{\mu - x}{\sigma^2},$$

so that f belongs to the Pearson family (3.4) with $d = -\mu$, $A_0 = \sigma^2$, and $A_1 = A_2 = 0$. It follows that

$$f(x) = \sigma^{-1}(2\pi)^{-1/2} \exp\left\{-\frac{(x-\mu)^2}{2\sigma^2}\right\},$$

a normal distribution with mean μ and variance σ^2 . Hence, (3.2) can be written as

$$E(X | X > t) = \mu + \sigma^2 r(t).$$

This is Theorem 3.1 of [13], see also [14,15].

3.2. The Maxwell Distribution

The Maxwell distribution has the form

$$f(x) = 4b^{-3}\pi^{-1/2}x^2 \exp\left\{-\frac{x^2}{b^2}\right\}, \quad x > 0, \quad b > 0,$$

see [16]. We then have

$$\frac{f'(x)}{f(x)} = \frac{2}{x} - \frac{2x}{b^2},$$

which can be written in form (3.1) with $\mu = 0$ and $g(x) = (1 + b^2/x^2)b^2/2$. Hence,

$$E(X | X > t) = \frac{(t^2 + b^2)b^2 r(t)}{2t^2}.$$

The fact that the corresponding failure rate is increasing can be seen by using Glaser's [17] result.

4. ULTIMATELY INCREASING FAILURE RATE DISTRIBUTIONS

Consider the class of distributions whose failure rate is ultimately increasing. More specifically, the failure rate should be strictly increasing from some point onward. Obviously, the important class of lifetime distributions having a bathtub-shaped failure rate with change points $0 \leq t_1 \leq t_2 < \infty$ (i.e., for which the failure rate is strictly decreasing on the interval $[0, t_1]$, constant on $[t_1, t_2]$ and strictly increasing on $[t_2, \infty)$) constitutes a proper subclass of the distributions we consider here.

We will see that if the failure rate is strictly increasing from some point onward, then under certain additional conditions the mean residual life can be expanded in terms of Gaussian probability functions.

NOTATION. Our conventions regarding the Bachmann-Landau O -notation, the Vinogradov \ll -notation, and o -notation are fairly standard. Thus, if h is a function of a positive real variable, the symbol $O(h(t))$, $t \rightarrow \infty$, denotes an unspecified function g for which there exist positive real numbers t_0 and B such that $|g(t)| \leq B|h(t)|$ for all real $t > t_0$. For such g , we write $g(t) \ll h(t)$ or $g(t) = O(h(t))$. The notation $g(t) = o(h(t))$, $t \rightarrow \infty$, means that for every real $\varepsilon > 0$, no matter how small, there exists a positive real number t_0 such that $|g(t)| \leq \varepsilon|h(t)|$ whenever $t > t_0$.

THEOREM 1. *Suppose that from some point onward, the failure rate r increases (strictly) without bound. Suppose further that for some positive integer n , the $n - 1$ derivative is continuous and satisfies*

$$\left| r^{(n-1)}(t+x) \right| \ll \left| r^{(n-1)}(t) \right|, \quad t \rightarrow \infty, \quad (4.1)$$

uniformly in x for $0 \leq x \leq \min(1, |r''(t)|^{-1/3})$, and

$$r^{(j)}(t) \ll \max\left(1, |r''(t)|^{(j+1)/3}\right), \quad t \rightarrow \infty, \quad 3 \leq j \leq n-1.$$

Finally, suppose there exists a positive real number ε such that for each integer j in the range $3 \leq j \leq n$,

$$r^{(j-1)}(t) = o\left((r(t))^{j-j\varepsilon}\right), \quad t \rightarrow \infty. \quad (4.2)$$

Then we have the following expansion for the mean residual life:

$$m(t) = \sum_{k=0}^{n-1} b_k(t) \varphi_k(t) + o\left((r(t))^{-1-n\varepsilon}\right), \quad t \rightarrow \infty, \quad (4.3)$$

where the coefficients $b_k(t)$ are given by the formal power series identity

$$\sum_{k=0}^{\infty} b_k x^k = \exp \left\{ - \sum_{k=3}^{\infty} r^{(k-1)}(t) \frac{x^k}{k!} \right\} \quad (4.4)$$

and

$$\begin{aligned} \varphi_k(t) &= \int_0^{\infty} x^k \exp \left\{ -xr(t) - \frac{1}{2}x^2 r'(t) \right\} dx \\ &= (-1)^k \sqrt{\frac{2\pi}{r'(t)}} \left[\frac{\partial^k}{\partial p^k} \left(1 - \Phi \left(\frac{p}{\sqrt{r'(t)}} \right) \right) \exp \left\{ \frac{p^2}{2r'(t)} \right\} \right]_{p=r(t)}. \end{aligned} \quad (4.5)$$

Here,

$$\Phi(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^x e^{-v^2/2} dv \quad (4.6)$$

is the Gaussian probability function, i.e., the cumulative distribution function of the standard normal distribution.

Before proving Theorem 1, we make some preliminary remarks and give two illustrative examples. First, if $r''(t) = 0$, then the uniformity condition on x in (4.1) should be interpreted as $0 \leq x \leq 1$. Next, observe that hypothesis (4.2) becomes more restrictive as ε increases. In particular, $\varepsilon > 1$ implies $\lim_{t \rightarrow \infty} r^{(j-1)}(t) = 0$. Of course, as ε increases, the error term in (4.3) decreases. On the other hand, if $0 < \varepsilon < 1$, then $r^{(j-1)}(t)$ does not necessarily approach zero, but the correspondingly weaker hypothesis implies a weaker conclusion (larger error term). In any case, since r increases without bound, the error term tends to zero as $t \rightarrow \infty$. Additionally, if r is infinitely differentiable, we may let $n \rightarrow \infty$ in (4.3) to obtain the *convergent* infinite series expansion

$$m(t) = \sum_{k=0}^{\infty} b_k(t) \varphi_k(t),$$

valid for all sufficiently large values of t . (More specifically, for those t for which $r(t) > 1$.)

In general, however, we do not assume the failure rate has infinitely many derivatives; n is fixed and the generating function (4.4) is a *formal* power series. Expanding (4.4) to compute $b_k(t)$ in terms of $r(t)$ and its derivatives shows that if r has only $n - 1$ derivatives, then $b_k(t)$ is undefined if $k \geq n$. Differentiating (4.4) leads to the recurrence

$$b_{k+1}(t) = -\frac{1}{k+1} \sum_{j=2}^k \frac{r^{(j)}(t)}{j!} b_{k-j}(t), \quad k \geq 2, \quad (4.7)$$

from which the coefficients $b_k(t)$ may be successively determined, starting with the initial values $b_0(t) = 1$, $b_1(t) = b_2(t) = 0$. On the other hand, an application of the multinomial theorem yields the explicit representation

$$b_k(t) = \sum_{p=0}^{\lfloor k/3 \rfloor} (-1)^p \sum \prod_{j \geq 2} \frac{1}{\alpha_j!} \left(\frac{r^{(j)}(t)}{(j+1)!} \right)^{\alpha_j}, \quad (4.8)$$

in which $\lfloor k/3 \rfloor$ is the greatest integer not exceeding $k/3$ and the inner sum is over all nonnegative integers $\alpha_2, \alpha_3, \dots$, such that $\sum_{j \geq 2} \alpha_j = p$ and $\sum_{j \geq 2} (j+1)\alpha_j = k$.

Finally, we note that the functions φ_k of (4.5) may also be given more explicitly. By setting $a = r(t)$ and $b = 2r'(t)$ in Lemma 1 below, we find that

$$\begin{aligned} \varphi_k(t) &= (-1)^k \left(\frac{2}{r'(t)} \right)^{(k+1)/2} \\ &\times \left\{ \sum_{h=0}^{\lfloor k/2 \rfloor} \binom{k}{2h} \lambda^{k/2-h} \Gamma\left(h + \frac{1}{2}\right) \left(\left(1 - \Phi(\sqrt{2\lambda})\right) e^\lambda + \frac{1}{2} \sum_{j=0}^{h-1} \frac{\lambda^{j+1/2}}{\Gamma(j+3/2)} \right) \right. \\ &\quad \left. - \frac{1}{2} \sum_{h=0}^{\lfloor k/2 \rfloor} h! \binom{k}{2h+1} \lambda^{(k-1)/2-h} \sum_{j=0}^h \frac{\lambda^j}{j!} \right\}, \end{aligned}$$

where $\lambda = (r(t))^2/2r'(t)$, $\Gamma(h+1/2) = \pi^{1/2} \prod_{j=1}^h (j-1/2)$, and Φ denotes the Gaussian probability function (4.6).

LEMMA 1. *Let a be a real number, let b be a positive real number, and let k be a nonnegative integer. Then,*

$$\begin{aligned}
 & \int_0^{\infty} x^k \exp \{-ax - bx^2\} dx \\
 = & (-1)^k b^{-(k+1)/2} \left\{ \sum_{h=0}^{\lfloor k/2 \rfloor} \binom{k}{2h} \lambda^{k/2-h} \Gamma \left(h + \frac{1}{2} \right) \left(\left(1 - \Phi \left(\sqrt{2\lambda} \right) \right) e^{\lambda} + \frac{1}{2} \sum_{j=0}^{h-1} \frac{\lambda^{j+1/2}}{\Gamma(j+3/2)} \right) \right. \\
 & \left. - \frac{1}{2} \sum_{h=0}^{\lfloor k/2 \rfloor} h! \binom{k}{2h+1} \lambda^{(k-1)/2-h} \sum_{j=0}^h \frac{\lambda^j}{j!} \right\},
 \end{aligned}$$

where $\lambda = a^2/4b$.

Proofs of Theorem 1 and Lemma 1 are relegated to Section 6, the final section.

5. APPLICATIONS

We provide two examples indicating how Theorem 1 may be applied.

EXAMPLE. Consider a linear failure rate of the form

$$r(t) = \alpha + \beta t, \quad \beta > 0. \quad (5.1)$$

The motivation and application of (5.1) to analyzing various data sets has been demonstrated by Kodlin [18] and Carbone *et al.* [19]. Statistical inference related to the linear failure rate model has been studied by Bain [20], Shaked [21], and more recently by Sen and Bhattacharya [22]. For this model, the hypotheses of Theorem 1 are trivially satisfied for *any* positive integer n and *any* positive real number ε . Since r'' vanishes identically in this case, we see that $b_k(t) = 0$ for $k > 0$ in (4.3) and in fact we have the exact result

$$m(t) = \int_0^{\infty} \exp \left\{ -(\alpha + \beta t)x - \frac{\beta x^2}{2} \right\} dx = \exp \left\{ \frac{(\alpha + \beta t)^2}{2\beta} \right\} \left(1 - \Phi \left(\frac{\alpha + \beta t}{\sqrt{\beta}} \right) \right) \sqrt{\frac{2\pi}{\beta}}.$$

EXAMPLE 2. Chen [23] proposes the two-parameter distribution with cumulative distribution function given by

$$F(t) = 1 - \exp \left\{ (1 - \exp(t^\beta)) \lambda \right\}, \quad t > 0,$$

where $\lambda > 0$ and $\beta > 0$ are parameters. The corresponding hazard function is the ultimately strictly increasing function of t given by

$$r(t) = \lambda \beta t^{\beta-1} \exp(t^\beta), \quad t > 0. \quad (5.2)$$

It is straightforward, albeit somewhat tedious, to verify that Chen's failure rate (5.2) satisfies the hypotheses of Theorem 1 with $n > 2$ and $0 < \varepsilon \leq 2/3$. Clearly, $\varepsilon = 2/3$ is optimal here. Thus, with derivatives of r in (4.4) and (4.5) now coming from (5.2), we see that the asymptotic formula

$$m(t) = \sum_{k=0}^{n-1} b_k \varphi_k(t) + o \left((r(t))^{-1-2n/3} \right), \quad t \rightarrow \infty,$$

holds for *all* integers $n > 2$. In particular, as the error term in (4.3) tends to zero in the limit as $n \rightarrow \infty$, we obtain the *convergent* infinite series representation

$$m(t) = \sum_{k=0}^{\infty} b_k(t) \varphi_k(t),$$

valid for all sufficiently large values of t . There is no need to work out the coefficients $b_k(t)$ explicitly in this case. One can simply use recurrence (4.7) to generate them.

6. PROOFS

PROOF OF THEOREM 1. Since the failure rate is strictly increasing from some point onward, there exists $t_0 \geq 0$ such that $r'(t) > 0$ for all $t \geq t_0$. Also, since $\lim_{t \rightarrow \infty} r(t) = \infty$, there exists $t_1 \geq 0$ such that $r(t) \geq 1$ for $t \geq t_1$. Now let $t \geq \max(t_0, t_1)$, $\delta = \delta(t) = \min(1, 1/\sqrt[3]{|r''(t)|})$, and set

$$I(t) := \int_0^\delta \exp\{R(t) - R(t+x)\} dx, \quad J(t) := \int_\delta^\infty \exp\{R(t) - R(t+x)\} dx,$$

so that $m(t) = I(t) + J(t)$. We have

$$\begin{aligned} J(t) &\leq \int_\delta^\infty r(t+x) \exp\{R(t) - R(t+x)\} dx \\ &= \exp\{R(t) - R(t+\delta)\} \\ &= \exp\left\{-\delta r(t) - \int_0^\delta x r'(t+\delta-x) dx\right\} \\ &\leq \exp\{-\delta r(t)\}. \end{aligned}$$

But $r'' = o(r^{3-3\varepsilon})$. By definition of δ , it follows that from some point onward, we must have $\delta r \geq \min(r, r^\varepsilon)$. Therefore, if we set $\nu = \min(1, \varepsilon)$, then $\nu > 0$ and

$$J(t) \leq \exp\{-(r(t))^\nu\}, \tag{6.1}$$

for all sufficiently large values of t .

Next, we write

$$\begin{aligned} I(t) &= \int_0^\delta \exp\left\{-\sum_{k=1}^{n-1} r^{(k-1)}(t) \frac{x^k}{k!} - \frac{1}{(n-1)!} \int_0^x u^{n-1} r^{(n-1)}(t+x-u) du\right\} dx \\ &= \int_0^\delta \left(\sum_{k=0}^{n-1} b_k(t) x^k + E_n(x, t)\right) \exp\left\{-xr(t) - \frac{1}{2}x^2 r'(t)\right\} dx, \end{aligned}$$

where

$$\begin{aligned} E_n(x, t) &= \exp\left\{-\sum_{k=3}^{n-1} r^{(k-1)}(t) \frac{x^k}{k!} - \frac{1}{(n-1)!} \int_0^x u^{n-1} r^{(n-1)}(t+x-u) du\right\} - \sum_{k=0}^{n-1} b_k(t) x^k \\ &= O\left(x^n \max \prod_{j=2}^{n-1} |r^{(j)}(t)|^{\alpha_j}\right), \end{aligned}$$

and the maximum is taken over all nonnegative integers α_j satisfying $\sum_{j=2}^{n-1} (j+1)\alpha_j = n$. In view of the fact that $r^{(j-1)} = o(r^{j-j\varepsilon})$ for $3 \leq j \leq n$, it follows that

$$E_n(x, t) = o(x^n (r(t))^{n-n\varepsilon}), \quad 0 \leq x \leq \delta.$$

If we now write

$$\begin{aligned} I(t) &= \sum_{k=0}^{n-1} b_k(t) \int_0^\infty x^k \exp\left\{-xr(t) - \frac{1}{2}x^2 r'(t)\right\} dx \\ &\quad - \sum_{k=0}^{n-1} b_k(t) \int_\delta^\infty x^k \exp\left\{-xr(t) - \frac{1}{2}x^2 r'(t)\right\} dx \\ &\quad + \int_0^\delta E_n(x, t) \exp\left\{-xr(t) - \frac{1}{2}x^2 r'(t)\right\} dx, \end{aligned}$$

then we find that

$$I = \sum_{k=0}^{n-1} b_k \varphi_k + \sum_{k=0}^{n-1} O(b_k r^k e^{-\delta r}) + o\left(r^{n-n\varepsilon} \int_0^\delta x^n e^{-xr} dx\right).$$

The hypotheses on r and the definition of the coefficients b_k imply that $b_k = O(r^{k-k\varepsilon})$. From the derivation of estimate (6.1) for J , we recall that $\exp(-\delta r) \leq \exp(-r^\nu)$, at least from some point onward. Finally, as

$$\int_0^\delta x^n e^{-xr} dx \leq \int_0^\infty x^n e^{-xr} dx = n! r^{-n-1},$$

it follows that

$$I = \sum_{k=0}^{n-1} b_k \varphi_k + o(r^{-1-n\varepsilon}). \quad (6.2)$$

Since $m = I + J$, combining (6.1) and (6.2) gives the stated result for m .

To complete the proof, it remains only to establish the asserted evaluation of the integrals φ_k . But this is readily obtained by completing the square in the exponential and differentiating under the integral. \blacksquare

PROOF OF LEMMA 1. By a straightforward change of variables, we find that

$$\begin{aligned} 2b^{(k+1)/2} \int_0^\infty x^k \exp\{-ax - bx^2\} dx &= e^\lambda \int_\lambda^\infty (t^{1/2} - \lambda^{1/2})^k e^{-t} t^{-1/2} dt \\ &= e^\lambda \sum_{h=0}^k \binom{k}{h} (-1)^{k-h} \lambda^{(k-h)/2} \Gamma\left(\frac{h+1}{2}, \lambda\right), \end{aligned} \quad (6.3)$$

where

$$\Gamma(\alpha, \lambda) = \int_\lambda^\infty t^{\alpha-1} e^{-t} dt \quad (6.4)$$

is the incomplete gamma function. If we integrate (6.4) by parts and then divide both sides of the result by $\Gamma(\alpha + 1) = \alpha\Gamma(\alpha)$, we obtain the recurrence formula

$$\frac{\Gamma(\alpha + 1, \lambda)}{\Gamma(\alpha + 1)} = \frac{\lambda^\alpha e^{-\lambda}}{\Gamma(\alpha)} + \frac{\Gamma(\alpha, \lambda)}{\Gamma(\alpha)},$$

which can be iterated to give

$$\frac{\Gamma(\alpha + k, \lambda)}{\Gamma(\alpha + k)} = e^{-\lambda} \sum_{h=0}^{k-1} \frac{\lambda^{\alpha+h}}{\Gamma(\alpha + h + 1)} + \frac{\Gamma(\alpha, \lambda)}{\Gamma(\alpha)}, \quad (6.5)$$

valid for any nonnegative integer k . In particular, when $\alpha = 0$,

$$\frac{\Gamma(k, \lambda)}{\Gamma(k)} = e^{-\lambda} \sum_{h=0}^{k-1} \frac{\lambda^h}{h!}. \quad (6.6)$$

Equation (6.6) is valid for all positive integers k if $\lambda \geq 0$; it is also valid when $k = 0$ if $\lambda > 0$.

Substituting $\alpha = 1/2$ in (6.5) yields

$$\begin{aligned} \frac{\Gamma(k + 1/2, \lambda)}{\Gamma(k + 1/2)} &= e^{-\lambda} \sum_{h=0}^{k-1} \frac{\lambda^{k+1/2}}{\Gamma(k + 3/2)} + \frac{\Gamma(1/2, \lambda)}{\Gamma(1/2)} \\ &= e^{-\lambda} \sum_{h=0}^{k-1} \frac{\lambda^{k+1/2}}{\Gamma(k + 3/2)} + 2\left(1 - \Phi(\sqrt{2\lambda})\right). \end{aligned} \quad (6.7)$$

Using (6.6) and (6.7), we get the stated result from (6.3). \blacksquare

REFERENCES

1. C. Bryson and M.M. Siddiqui, Some criteria for aging, *Journal of the American Statistical Association* **64**, 1472–1483, (1969).
2. W. Hollander and F. Proschan, Tests for mean residual life, *Biometrika* **62**, 585–593, (1975).
3. E.J. Muth, Reliability models with positive memory derived from the mean residual life function, In *Theory and Applications of Reliability*, (Edited by C.P. Tsokos and I.N. Shimi), pp. 401–434, Academic Press, (1977).
4. I. Meilijson, Limiting properties of the mean residual lifetime function, *Annals of Mathematical Statistics* **43** (1), 354–357, (1972).
5. A.A. Balkema and L. de Hann, Residual life time at great age, *Annals of Probability* **2** (5), 792–804, (1974).
6. D.M. Bradley and R.C. Gupta, Limiting behaviour of the mean residual life, *Annals of the Institute of Statistical Mathematics* (to appear).
7. Y.P. Chaubey and P.K. Sen, On smooth estimation of mean residual life, *Journal of Statistical Planning and Inference* **75**, 223–236, (1999).
8. J.M. Ruiz and J. Navarro, Characterization of distributions by relationships between failure rate and mean residual life, *IEEE Transactions on Reliability* **43** (4), 640–644, (1994).
9. U.N. Nair and P.G. Sankaran, Characterization of the Pearson family of distributions, *IEEE Transactions on Reliability* **40** (1), 75–77, (1991).
10. A.N. Ahmed, Characterization of beta, binomial and Poisson distributions, *IEEE Transactions on Reliability* **40** (3), 290–295, (1991).
11. S. Osaki and X. Li, Characterizations of gamma and negative binomial distributions, *IEEE Transactions on Reliability* **37** (4), 379–382, (1988).
12. S.M. El-Arshy, Useful relationships between the log-survival function and truncated moments, with applications, *Statistical Papers* **36**, 145–154, (1995).
13. A.N. Ahmed and A.A. Abdul-Rahman, On characterization of the normal, one-sided normal, lognormal, and logistic distributions via conditional expectations, *Pakistan Journal of Statistics* **9** (3)B, 19–30, (1933).
14. J.I. McGill, The multivariate hazard gradient and moments of the truncated multinormal distribution, *Communications in Statistics Theory and Methods* **21** (11), 3053–3060, (1992).
15. S. Kotz and D.N. Shanbhag, Some new approaches to probability distributions, *Advances in Applied Probability* **12**, 903–921, (1980).
16. S.M. El-Arshy, Characterizations of Weibull, Rayleigh and Maxwell distributions using truncated moments and order statistics, *Egyptian Journal of Statistics* **37** (2), 198–212, (1993).
17. R.E. Glaser, Bathtub and related failure rate characterizations, *Journal of the American Statistical Association* **75**, 667–672, (1980).
18. D. Kodlin, A new response time distribution, *Biometrics* **2**, 227–239, (1967).
19. P.O. Carbone, L.E. Kellerhouse and E.A. Gehan, Plasmacytic myeloma: A study of relationship of survival to various clinical manifestations and anomalous protein type in 112 patients, *American Journal of Medicine* **42**, 937–948, (1967).
20. L.J. Bain, Analysis for the linear failure rate life testing distribution, *Technometrics* **16** (4), 551–559, (1974).
21. M. Shaked, Accelerated life testing for a class of linear hazard rate type distributions, *Technometrics* **20** (4), 457–466, (1978).
22. A. Sen and G.K. Bhattacharya, Inference procedures for the linear failure rate model, *Journal of Statistical Planning and Inference* **46**, 59–76, (1995).
23. Z. Chen, A new two-parameter lifetime distribution with bathtub shape or increasing failure rate function, *Statistics & Probability Letters* **49**, 155–161, (2000).